



Keywords

Dehydration,
Radical Protrusion,
Root Hair

Received: June 21, 2017

Accepted: July 5, 2017

Published: October 17, 2017

Mechanisms of Desiccation Tolerance in Plants during Seed to Seedling Transition

Adane Gebeyehu

National Agricultural Biotechnology Research Center, Ethiopia

Email address

adyamrot@gmail.com

Citation

Adane Gebeyehu. Mechanisms of Desiccation Tolerance in Plants during Seed to Seedling Transition. *International Journal of Agricultural and Biosystems Engineering*.

Vol. 2, No. 5, 2017, pp. 34-43.

Abstract

The present study was conducted to characterize the role of *Arabidopsis thaliana* genes in desiccation tolerance during seed to seedling transition. Dehydration stress, such as drought and desiccation, affects the physiological and biochemical activities of seeds and seedlings through either suppressing or inducing stress responsive genes of plants. Responsiveness of genes related to desiccation tolerance (DT) can be mediated with increase in abscisic acid (ABA) level. Treatment of plants with exogenous ABA, therefore, induces responses of stress-associated genes. The objective of this study was to characterize *Arabidopsis thaliana* stress-associated genes for their role in DT in germinated *A. thaliana* seeds. Four different mutant lines of *Arabidopsis* (*sod1*, *daa1*, *di19* and *vip1*) were used, which stress-associated genes are knocked out. Seeds of these mutants were selected at two stages (RP-Radical Protrusion and RH-Root Hair), and analyzed for ability to acquire DT in mild-osmotic stress by applying PEG. Levels of gene expression were analyzed using RT-qPCR at RP and RH before and after PEG treatment. Eight key genes of the ABA response pathway were selected and checked for their expression level. Quantitative gene expression data for RP and RH stages showed an overall trend to increase the expression level after PEG treatment. Expression of genes of ABA biosynthesis and the receptors increase in response to PEG treatment, while ABA catabolic gene decreases. The author conclude that ABA genes are assumed to be involved in response to DT, although, three mutants: *sod1*, *daa1* and *di19* showed an increase of 20% in acquisition of DT compared to the wild type, genes involved in ABA pathway did not show difference related to phenotype.

1. Introduction

The seed is an important dispersal unit in the plant life cycle with respect to its survival as a species. It can survive the period between seed maturation and seedling establishment. For successful survival, seeds should tolerate adverse environmental conditions such as drought, cold, salinity, low or high temperature as well as heavy metals (Bray et al., 2000; Krasensky and Jonak, 2012; Yang and Chen, 2013). Drought affects the physiological and biochemical activities of seeds through inducing stress-related gene expression (Yamaguchi-Shinozaki & Shinozaki, 2005; Shinozaki, K., & Yamaguchi-Shinozaki, K., 2007). For this reason, traits such as desiccation tolerance and seed dormancy are very important. According to the IWMI report, one-third of the world's population (~1 billion) will live in water deficit areas by the year 2025 (www.iwmi.org). This will result in severe economic impact on the agricultural sector. For this reason, it is vital to better understand stress tolerance mechanisms and develop

new tools to engineer plants that are better adapted to water scarcity in these areas for better access to food and raw materials to support the agricultural sector.

There are two types of tolerance based on the critical water level. These are drought tolerance: the capacity to tolerate moderate dehydration, to moisture content below $\sim 0.3\text{g H}_2\text{O}$ per gram dry weight (Hoekstra et al., 2001; Tripathi, et al., 2014). And desiccation tolerance or anhydrobiosis: the tolerance of further dehydration, to water levels below $0.1\text{g H}_2\text{O}$ per gram of dry weight and successive re-hydration without permanent damage in periods of long dry periods (Hoekstra et al., 2001; Maia et al., 2011; Dekkers et al., 2015).




Figure 1. Biosynthesis of ABA from carotenoid precursors in higher plants.

Recently, varieties of molecular and genomic level research have been conducted in the model plant *Arabidopsis thaliana* (Dekkers et al., 2015). Stress-inducible genes, proteins and metabolites involved in desiccation tolerance are identified using microarray analysis (Shinozaki et al., 2003; Todaka, Daisuke, et al., 2012; Sarhadi et al., 2012; Maia et al., 2011). Most of the stress-inducible genes are found to be involved in the synthesis of osmolytes and enzymes that are

involved in ABA biosynthesis (Bartels & Sunkar, 2005). ABA biosynthetic genes and enzymes have been identified in *Arabidopsis thaliana*. For example, zeaxanthin epoxidase (ZEP) is one of these enzymes, which is encoded by the ABA1 gene of *Arabidopsis thaliana* (Xiong et al. 2002; Xiong & Zhu 2003; Barerro et al., 2006). It catalyzes a step of the ABA biosynthetic pathway (Figure 1) through catalyzing epoxidation of zeaxanthin to antheraxanthin and all-trans violaxanthin. All-trans-violaxanthin is then converted into the cis-neoxanthin and cis violaxanthin, which are cleaved by a 9-cis epoxy carotenoid dioxygenase (NCED) to form xanthoxin (Iuchi et al. 2000). Then xanthoxin is converted to abscisic aldehyde by the catalyzing ABA2 enzyme of *Arabidopsis thaliana*. The abscisic aldehyde is then oxidized to form ABA by the enzyme aldehyde oxidase 3 (AAO3). As a result, ABA is derived from the oxidative cleavage of a 9-cis-epoxy-carotenoid (Schwartz & Zeevaart, 2010).

DT is a complex trait that requires activities involved in abscisic acid (ABA) signaling. There are three core components that are involved in ABA signaling: PYR/PYL/RCAR receptor family consisting of pyrabactin resistance 1 (PYR1)-like regulatory components of ABA receptors, the type 2C protein phosphatases (PP2Cs) and the sucrose-non-fermenting kinase 1-related protein kinase 2 (SnRK2s) family (Umezawa et al., 2010; Komatsu et al., 2013). PP2C phosphatases function as a negative regulator of ABA signaling components (Cutler et al., 2010) whereas the SnRK2s are positive regulator of ABA signaling (Umezawa et al., 2010). This indicates that signaling complex of these three protein families are important to regulate ABA response in plants. Under normal conditions, PP2C inactivates SnRK2 by dephosphorylating its multiple residues to repress the ABA signaling but when endogenous ABA is up regulated due to abiotic stress, the PYR/PYL/RCAR binds to ABA and interacts with PP2Cs to inhibit protein phosphatase activity. PYL proteins are therefore the most important to inhibit ABA-dependent PP2C phosphatase activity (Ma et al. 2008). The closed gate loop of PYLs is very important to seal the catalytic site of PP2Cs and expose the serine residue, an interface that interacts with group A PP2C upon ABA binding. Then, the ABA-bound receptor inhibits the PP2C phosphatase activity suggesting that PYLs are negative regulatory sub units of PPCs during ABA signaling (Ma et al., 2009). When the PP2C activity is inhibited, SnRK2 are released from PP2Cs-dependent regulation and turn to ABA signaling by phosphorylating the downstream basic leucine zipper (bZIP)-type transcription factors such as ABA response elements (ABRE)/ABRE binding factors (ABF) or membrane proteins (Umezawa et al., 2010).

In addition, five ABA INSENSITIVE (ABI1 to ABI5) loci that are involved in ABA downstream signaling are identified in *Arabidopsis*. They affect *Arabidopsis* germination and post-germination processes (Finkelstein, 2013). The ABI1 and ABI2 genes encode homologous type 2C protein Ser/Thr

Phosphatases (PP2Cs). The remaining three ABA Insensitive genes, ABI3, ABI4 and ABI5 encode transcription factors of the basic B3-domain, APETALA2-domain (AP2) and basic leucine zipper-(bZIP) domain families respectively (Lopez-Molina & Chua, 2000). They act as transcription factors found to be abundantly expressed in seeds and depending on the gene targeted; they act as either activators or repressors (Finkelstein, 2013). The ABA Insensitive genes; ABI3, ABI4 and ABI5, regulate genes that are involved in many physiological processes such as lipid metabolism, postembryonic development, photosynthesis, response to ABA stimulus and chlorophyll metabolic processes (Wind et al., 2013). The ABA insensitive gene ABI4 is essential for regulation of ABA and sugar signaling during seed germination and early seedling development (Huijser et al., 2000). It also activates expression of its own gene, ABI5 and starch binding enzyme SBE2.2 through binding directly to the promoter region of all three genes (Huijser et al., 2000). Dry seeds of *A. thaliana* *abi5* mutants showed lower levels of ABRE than the wild type suggesting that ABI5 has an important role in response to exogenous ABA during germination by protecting seeds from water loss during drought stress (Lopez-Molina et al., 2001).

Several studies have been conducted to explore the molecular basis of gain, loss and re-establishment of desiccation tolerance in germinated *abi* mutants of *Arabidopsis* (Maia et al., 2011; Verdier et al., 2013). Studies revealed that wild type Columbia (Col-0) seeds lose desiccation tolerance during germination and can re-acquire it during germination by application of a mild osmotic stress (Maia et al., 2011). This has been proposed to be a good method to investigate the molecular basis of acquisition of desiccation tolerance. Previous studies have shown that over expression of stress-inducible genes that accumulate during the acquisition of desiccation tolerance are useful to improve it (Sun et al., 2013). Therefore, the objective of this study was to find out which genes were involved in desiccation tolerance in two germination stages (RP and RH) of *Arabidopsis*. Four different mutant lines of *Arabidopsis* were used, which stress-inducible genes are knocked out. The *vip1* showed greatest result, where each gene related to ABA biosynthesis and receptors showed a decreased trend between treated and untreated stages. These results hypothesize that ABA biosynthesis and signaling are associated to its phenotype, which did not acquire DT at RH stage.

Table 2. Knock-out mutants used for evaluation of desiccation tolerance (<http://www.arabidopsis.org>).

| Gene | Function | Other Names | Description |
|-----------|-----------------------------|-----------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| AT1G12520 | Copper chaperone | ATCCS, CCS, Copper chaperone for SOD1 | Copper-zinc superoxide dismutase copper chaperone (delivers copper to the Cu-Zn superoxide dismutase). Up-regulated in response to copper and senescence. |
| AT1G64110 | Coding for ATPase | DUO1-Activated ATPase1 (DAA1) | Target promoter of the male germline-specific transcription factor DUO1 |
| AT1G56280 | Drought induced gene family | ATDI19, Drought-Induced 19 (DI19) | Encodes a gene whose transcript level in root and leaves increases to progressive drought stress. The increase in transcript level is independent from abscisic acid level. It appears to be a member of plant-specific gene family. |
| AT3G01310 | Phosphate metabolism | <i>Arabidopsis</i> Homolog of Yeast VIP1 1 (ATVIP1) | Encodes a myoinositol hexakisphosphate kinase. |

2. Materials and Methods

Genotyping the *Arabidopsis thaliana* mutant lines DNA Extraction

DNA extraction was performed with unexpanded young leaves of the knockout mutants for selection of homozygous T-DNA lines. The tissue material was grinded into non-sterile polypropylene 1.2 ml tubes (Collection Micro tubes (racked, 8 x 12) having one metal bead inside and 300µl extraction buffer (Table 1). Then the samples were shaken for 1 minute at 30Hz, and incubated in an oven at 600C for 1 hour. Afterwards, samples were centrifuged at 3920 rpm for 15 minutes and 75µl of the upper watery phase of each sample (supernatant) was pipetted into new microplates. Then 75µl isopropanol and 30µl 10mM NH₄AC were added to each sample in order to precipitate the nucleic acids. Samples were kept for 15 minutes at room temperature. Then the samples were centrifuged at 3920 rpm for 15 minutes. The supernatant was poured off and 50µl 70% ethanol was added; centrifuged at 3920 rpm for 5 minutes to wash the pellet and ethanol was poured off. The pellets were dried at room temperature for 30 minutes and re-suspend in 50µl MQ water. Quality of the DNA was checked by Nanodrp®.

Table 1. Extraction buffer preparation.

| Extraction buffer stock | for 500 ml: |
|-------------------------|--------------------------|
| 0.35 M Sorbitol | 31.9 g Sorbitol |
| 0.1 M Tris-HCl pH 8.0 | 50 ml 1M Tris-HCl pH 8.0 |
| 5 mM EDTA pH 8.0 | 5 ml 0.5M EDTA pH 8.0 |

PCR Assays

In order to investigate the presence or absence of the T-DNA insertion, PCR analysis with the border primers SAIL_534_B01 (for the mutant AT1G33970) & SALK for the other nine mutant lines and primer pairs of the respective mutants were performed. Gel electrophoresis was used to check the DNA samples and the corresponding wild type control for homozygosity. Therefore, four Knock-out mutant lines of *Arabidopsis thaliana* that were previously identified by Barrero et al., (2010); Cabello et al., (2012) and Ribone et al., (2015) has been homozygous were used for further analysis.

Phenotyping the *Arabidopsis thaliana* mutant lines

Seeds of the wild type accession, Columbia (Col-0) and four knockout mutant lines of *Arabidopsis thaliana* (Table 2) have been used for this study.

First, seeds of the above mentioned four knockout mutant lines of *Arabidopsis thaliana* and the accession Columbia

(Col-0) were put in 9 cm Petri dishes on two layers of blue filter paper (Anchor paper Co.) moistened with 10 ml demi-water. Then, seeds were put in a dark room to be stratified at 4°C for 72 hours to break dormancy. After stratification, the seeds were transferred to an incubator with constant white light at 22°C and allowed to germinate. After approximately 28 hours (RP) and 32 hours (RH) (Figure 2), the germinated seeds were selected at the specific stages by hand using a stereomicroscope. Three hundred seeds of each biological replicate for each stage were selected and 150 seeds were stored in -80°C and 150 seeds were used for treated with polyethylene glycol (PEG 8000 with a concentration of -2.5MPa).




Figure 2. Seed-to-seedling developmental stages of *Arabidopsis* seeds.

PEG Treatment

Many studies have been conducted on the attainment of desiccation tolerance during seed development and on its loss upon germination. In order to re-establish DT in the seeds of the 5 tested mutants (*sod1*, *daa1*, *di19* and *vip1*), a PEG treatment was performed as described by Maia et al., (2011) with slight modifications. A total number of 150 germinated seeds each for the RP and the RH stage for each line and replica were selected and incubated for three days in a PEG8000 solution with an osmotic potential of -2.5MPa. After incubation, seeds were washed with distilled water to remove the remaining PEG and kept at -80°C.

RNA Extraction

RNA extraction was performed using 150 germinated seeds of RP and RH stages after PEG incubation and 150 non-treated germinated seeds at the same developmental stages. Seeds were put into 2ml tubes and stored at -80°C prior to grinding. Total RNA was extracted according to the hot borate protocol (Wan and Wilkins, 1994). The RNA pellets were dissolved in 20µl DEPC MQ water and the RNA quality was checked by agarose gel electrophoresis. The absorbance (A260, 280 and 230) was measured using a Nanodrop® to check the quantity and quality of RNA.

cDNA Synthesis

cDNA was synthesized using the iScript™ cDNA synthesis kit (Bio-Rad) protocol. First, 5µl of master mix which contained 4µl 5x iScript reaction buffer and 1µl iScript reverse transcriptase was added to each tube. Then, 1µg of each RNA sample and MQ water was added to a final reaction volume of 20µl. The reaction mix was incubated at 25°C for 5 minutes, 42°C for 30 minutes, 85°C for 5 minutes

and subsequently kept at 4°C. Finally, one tenth of the cDNA reaction volume was diluted 10x with sterile MQ water followed by RT-qPCR using gene-specific primers.

Real-time quantitative PCR (RT-qPCR) Analysis

Real-time quantitative PCR (RT-qPCR) was performed on CFX96 connect TM Real-Time PCR Detection System (Bio-Rad). Amplification was done in a 10µl reaction mixture containing, 5µl SYBR Green, 1µl of 10µM primers, 2.5µl cDNA and 1.5 µl MQ water. Primer pairs were designed using the CLC Main Workbench software. The RT-qPCR program run with the first step at 95°C for 3 minutes, followed by 40 cycles at 95°C for 10 seconds and 60°C for 30 seconds followed by a melt curve 55°C to 95°C; increasing 0.5°C for every 5 seconds. Data were analyzed using CFX Data Analysis Manager Software and qbase+®. Primer efficiencies were measured with LinRegPCR software and most of the primers showed efficiencies higher than 90%. Melting curves were analyzed and a single peak was observed for all primers, confirming the specificity of the primers. Four reference genes identified by Dekkers et al. (2012) (*At1g16970*, *At2g43770*, *At3g12210*, and *At4g02080*) were tested and their expression stability was calculated by geNorm® software. Normalization of reference genes was done according Dekkers et al, (2012). Finally, calibrated normalized relative quantity (CNRQ) values were exported from qbase+® and statistically analyzed.

3. Results

Assessment of DT re-establishment in germinated *Arabidopsis thaliana* mutant lines




Figure 3. Response of mutant lines in acquisition of DT.

Four knock out mutants and the wild type Col-0 were selected according to their response in acquisition of DT (Figure 3) to check if ABA pathway genes are associated to DT in germinated seeds after PEG treatment at RP and RH stages. Mutants were *At1g12520* gene that codes for copper chaperone (*sod1*), *At1g64110* codes for ATPase (*daa1*), *At1g56280* codes for drought induced gene family member (*di19*) and *at3g01310* codes for a phosphoglycerate mutase-like gene (*vip1*). The superoxide dismutase (SOD) enzyme catalyzes the dismutation of superoxide radicals and involved in defense against Reactive oxygen species (Beyer et al., 1991). The SOD1 gene is shown up regulated in response to

No ABA biosynthesis genes showed significant changes in their expression levels after PEG treatment when compared to untreated seeds in both stages of development (Figure 4 a, b, c, d, e). The expression of NCED3 and NCED5 did not increase significantly with PEG treatment. Generally, there was a trend to increase in expression level after PEG treatment in contrast to the untreated seeds. These might suggest that, ABA content increases in response to PEG treatment. Gene expression is shown for untreated seeds and seeds treated for 3 days in PEG (-2.5 MPa). Bars represent the mean values \pm SE of three independent biological replicates. Small letters represent statistical differences ($p \leq 0.05$) between different developmental stages and different treatments. Bars with the same letter(s) are not significantly different at $p < 0.05$.




Figure 4. Gene expression (CNRO) for ABA biosynthesis genes *ABA1*, *NCED3*, *NCED5*, *NCED6* and *NCED9*.

ABA catabolic gene

The ABA catabolic gene, CYP707A2 does not show significant changes in its expression level after PEG treatment when compared to untreated seeds in both stages of development (Figure 5). However, there was a tendency to decrease in expression level after PEG treatment in contrast to the untreated seeds. This might suggest that, the CYP707A2 gene degrades the ABA content in PEG treated

seeds and as a result, PEG-treated seeds might not have accumulated ABA compared to untreated seeds. Gene expression is shown for untreated seeds and seeds treated for 3 days in PEG (-2.5 MPa). Bars represent the mean values \pm SE of three independent biological replicates. Small letters represent statistical differences ($p \leq 0.05$) between different developmental stages and different treatments. Bars with the same letter(s) are not significantly different at $p < 0.05$.




Figure 5. Gene expression analysis by RT-qPCR represented as CNRQ for ABA catabolic gene CYP707A2.

ABA receptors from the PYL/PYR/RCAR family

The expression levels of PYL7 and PYL9 did not show significant changes after PEG treatment when compared to untreated seeds in both stages of development (Figure 6). Generally, there was a trend to increase in expression level after PEG treatment in contrast to the untreated seeds. This might suggest that these genes are involved in DT regulation in response to osmotic stress and increased ABA sensitivity in germinating seeds.




Figure 6. Gene expression analysis (CNRQ) for ABA receptors from the PYL/PYR/RCAR family.

Gene expression is shown for untreated seeds and seeds treated for 3 days in PEG (-2.5 MPa). Bars represent the mean values \pm SE of three independent biological replicates. Small letters represent statistical differences ($p \leq 0.05$) between different developmental stages and different treatments. Bars with the same letter(s) are not significantly different at $p < 0.05$.

4. Discussion

Genotyping the Arabidopsis thaliana mutant lines

It is possible to detect the presence of T-DNA insertion of a given gene in Arabidopsis knock-out mutants by PCR using one T-DNA specific primer and gene specific primers to the respective knock-out mutant. As a result, the T-DNA is landed to the gene of interest and a PCR product is formed through amplification process. The T-DNA insert to be found within a gene is determined by genome size, gene size and number of T-DNA inserts in the population (Krysan et al., 1999). Since Arabidopsis thaliana is reproduced by self-fertilization, heterozygosity is rarely a problem. Therefore, all of the tested mutant lines were heterozygous for the T-DNA insertions. This might be due to disruption of gene function, homozygous lethal mutations or additional segregation of the T-DNA insertion into different genes. For example, 5 to 25kb long T-DNA insert can cause disruption of gene function because of very little intergenic material and small introns in Arabidopsis plants (Krysan et al., 1999). Due to this disruption of gene function, homozygous lethal mutations can happen in the form of heterozygous plants (Krysan et al., 1999). It might also be because of endogenous interference or silencing of the T-DNA insert.

Gene expression analysis of the ABA Pathway

ABA is known to be involved in regulating plant physiological processes such as seed dormancy. Seed dormancy can be influenced by altered ABA biosynthesis even though ABA deficient and ABA insensitive (ABI) mutants can produce desiccation tolerant (DT) seeds during maturation (Cutler et al., 2010). However, DT was shown to be lost during germination, on which the *Arabidopsis thaliana* seeds become sensitive to desiccation (Maia et al., 2011). Then, DT can be re-established when mild osmotic stress (like for example PEG) is applied to germinated seed up to radicle protrusion (RP), when seeds are at root hair (RH) stage acquisition of DT drops dramatically (Maia et al., 2011). The mechanisms involved how a seed or seedling perceives this osmotic signal is not clear yet. Therefore, knowing the key genes involved in ABA biosynthesis, signaling and degradation is very important.

ABA biosynthesis genes

The level of ABA biosynthesis and ABA sensitivity determines the response of plants to ABA. These propose that, increased ABA level triggers ABA mediated stress response in desiccation sensitive *Arabidopsis* seeds through the application of mild osmotic stress (PEG) (Maia et al, 2014). However, our data revealed that no ABA biosynthesis genes showed significant changes in their expression levels at both stages of development even though there was a trend to increase in expression levels after PEG treatment. These suggest that, ABA content increases in response to PEG treatment. The present result was in agreement with Maia et al 2014, on which all ABA biosynthesis genes showed a tendency to increase in their expression levels upon PEG treatment. In addition, DT can be rescued by PEG treatment regardless of increased ABA levels; since all NCEDs increased in their level of expression in PEG treated *Arabidopsis* seeds. According to a report by Westwood et al., (2013), plants expressing cucumber mosaic virus (CMV) factors showed increased drought tolerance without over accumulation of ABA. This suggests that, adjustment of ABA signalling and/or perception can be sufficient to induce a proper stress response besides enhancing ABA levels since increased ABA level is not always necessary to elicit DT (Maia et al, 2014). One of the reasons why ABA biosynthesis genes are not up regulated in response to PEG treatment might be due to the presence of the ABA catabolic genes such as the CYP707A family (Okamoto et al., 2006); inhibitors of ABA biosynthesis (Debeaujon a& Koornneef, 2000) and/or ABA conjugation products (Lim et al., 2005). It might also be due to the balance between ABA biosynthesis and ABA degradation. Furthermore, it might also be due to ABA translocation from one sub-organelle to the other. For example, NCEDs are found in the plastid but they differ in their binding to the thylakoid membrane. NCED2, NCED3 and NCED6 are localized in both stroma and thylakoid. Whereas, NCED9 is localized only in the stroma and NCED5 is bound by the thylakoid membrane. This difference in localization might affect the release of xanthoxin to the cytoplasm and some enzymatic activities during ABA biosynthesis.

ABA catabolic gene

ABA levels can also be regulated through catabolism in

addition to ABA biosynthesis. ABA catabolism has two pathways namely hydroxylation of ABA by the P-450 type monooxygenases to give unstable 8'-OH-ABA and esterification/conjugation of ABA to ABA-GE (Lim et al., 2005). The ABA-8'-hydroxylases are encoded by the CYP707A family having different patterns of expression in both time and space but all induced by both dehydration and subsequent rehydration. Because of spatial and temporal differences in expression patterns, each member of this family has different developmental and/or physiological role(s) (Okamoto et al., 2006). For example, CYP707A1 and CYP707A3 have a role for ABA catabolism in the middle of seed development but, CYP707A2 during imbibition and at the end of seed development (Okamoto et al., 2006). It has been shown that, CYP707A mutants in *Arabidopsis* accumulate more amount of ABA than lines over expressing ABA biosynthetic enzymes (Umezawa et al. 2006). In addition, overexpression of CYP707A effectively reduces endogenous ABA (Okamoto et al. 2006, Umezawa et al. 2006). As a result, CYP707As has a role in controlling the level of ABA biosynthesis. However, our data shown that the ABA catabolic gene, CYP707A2 does not show significant changes in its expression level after PEG treatment when compared to untreated seeds in both stages of development. Nevertheless, there were lines with a tendency to increase in expression level after PEG treatment. For example, the At1g56280 gene knocked out mutant line that code for drought induced gene family (di19) and the At3g01310 gene knocked out mutant line that codes for a phosphoglycerate mutase-like gene (vip1) showed to increase in level of expression up on PEG treatment. The reason for down regulation of the ABA catabolic gene, CYP707A2 in response to PEG treatment might be due to the fact that, the CYP707A2 gene degrades the ABA content in PEG treated seeds; as a result, PEG-treated seeds did not accumulate active ABA when compared to untreated seeds. It might also be due to continuous balance between ABA biosynthesis and ABA catabolism or these products were not detected by the PEG treated seeds due to some catabolic inhibitors that targets CYP707A. Whereas, the di19 mutant that has dehydration-induced and ABA-independent DT mechanism but not through ABA catabolism since it was ABA-independent. In addition, the vip1 mutant that encodes InsPs has a role in phosphate metabolism by controlling the anti-oxidant tocopherol, stress responses and hormonal signaling through providing a chemical code for plants to respond to these signals including signals of ABA catabolism. Due to these reasons, both mutants were shown up-regulated up on PEG treatment through detecting signals of ABA catabolism.

ABA receptors from the PYL/PYL/RCAR family

Three core components that are involved in ABA signal transduction pathways have been identified after intensive study on ABA (Miyakawa et al., 2013). These are PP2Cs, PYR/PYL/RCARs and SnRK2s (Umezawa et al., 2010; Komatsu et al., 2013). The PYR/PYL/RCAR receptor consists of pyrabactin resistance 1 (PYR1)-like regulatory components of ABA receptors that inhibits protein phosphatase activity

through inactivating PP2Cs (Ma et al., 2009). The PP2C phosphatases function as a negative regulator of ABA signaling components through inactivating SnRK2s (Cutler et al., 2010) whereas the SnRK2s are positive regulator of ABA signaling (Umezawa et al., 2010). Under a normal condition, PP2C inactivates SnRK2 by dephosphorylating its multiple residues but when endogenous ABA is up regulated due to abiotic stress such as desiccation and drought, the PYR/PYL/RCAR binds to ABA and interacts with PP2C to inhibit protein phosphatase activity. According to Gonzalez-Guzman et al., (2012), the PYR/PYL/RCAR protein family except PYL13 is capable of activating ABA-responsive gene expression that confirms this ABA receptors other than PYL13 has a role to regulate ABA signaling in response to PEG treatment in germinated seeds. However, our data revealed that no ABA receptors from the PYL/PYR/RCAR family showed significant changes in their expression levels at both stages of development even though there was a fashion to increase in expression levels after PEG treatment. The present result was in contrast with Maia et al., 2014, who found increased expression of PYL7 and PYL9 in response to PEG treatment in both RP and RH stages. The reason for non-significant changes in expression levels after PEG treatment might be due to conformational changes during ABA binding to the PYR/PYL/RCARs or different receptor proteins are involved in the ABA signaling pathway since some PYR/PYL proteins such as PYL9 and PYL5 have preference on binding to different ABA arrangement (Santiago et al., 2009). Therefore, analysis that is more careful has to be repeated to confirm the hypothesis.

Abbreviations

ABA-Absciscic acid
 CMV -Cucumber mosaic virus
 DT -Desiccation Tolerance
 IWMI -International Water Management Institute
 PEG-Poly Ethylene Glycol
 RH-Root Hair
 RP-Radical Protrusion and
 RT-qPCR-Real Time quantitative PCR

Authors' Contributions

The sole Author contributed in all stages of the experiment including final approval of manuscript.

Acknowledgements

This work was supported by the National Agricultural Biotechnology Research Center, Ethiopia. The Author would like to thank AASCIT for waiving the article processing charge.

Competing Interests

The author declare that no competing interest being the

sole Author.

Ethical Approval

All applicable institutional guidelines for the care and use of knocked out mutants were followed. This article does not contain any studies with human and animals participants performed by the author.

Funding

This study was funded by the EIAR-National Agricultural Biotechnology Research Center, Ethiopia.

Authors' Affiliation

Researcher, EIAR-National Agricultural Biotechnology Research Center, Ethiopia.

References

- [1] Abdel-Ghany SE, Müller-Moulé P, Niyogi KK, Pilon M, Shikanai T (2005) Two P-type ATPases are required for copper delivery in *Arabidopsis thaliana* chloroplasts. *Plant Cell*, 7: 1233–1251.
- [2] Ali, M. A., Plattner, S., Radakovic, Z., Wieczorek, K., Elashry, A., Grundler, F. M., & Bohlmann, H. (2013). An *Arabidopsis* ATPase gene involved in nematode - induced syncytium development and abiotic stress responses. *The Plant Journal*, 74 (5), 852-866.
- [3] Barker, C. J., Illies, C., Gaboardi, G. C. and Berggren, P. O. (2009) Inositol pyrophosphates: structure, enzymology and function. *Cell. Mol. Life Sci.* 66, 3851–3871.
- [4] Barrero JM, Millar AA, Griffiths J, Czechowski T, Scheible WR, Udvardi M, Reid JB, Ross JJ, Jacobsen JV, Gubler F. (2010) Gene expression profiling identifies two regulatory genes controlling dormancy and ABA sensitivity in *Arabidopsis* seeds. *Plant J* 61: 611–622.
- [5] Barrero, J., Rodríguez, P. L., Quesada, V., Piqueras, P., Ponce, M. R., & Micol, J. L. (2006). Both abscisic acid (ABA)-dependent and ABA - independent pathways govern the induction of NCED3, AAO3 and ABA1 in response to salt stress. *Plant, cell & environment*, 29 (10), 2000-2008.
- [6] Bartels, D., & Sunkar, R. (2005). Drought and salt tolerance in plants. *Critical reviews in plant sciences*, 24 (1), 23-58.
- [7] Beyer W, Imlay J, Fridovich I (1991). Superoxide dismutases. *Program for nucleic acid research in molecular biology*, 40: 221–253.
- [8] Borg, M., Brownfield, L., Khatab, H., Sidorova, A., Lingaya, M. and Twell, D. (2011). The R2R3 MYB transcription factor DUO1 activates a male germline-specific regulation essential for sperm cell differentiation in *Arabidopsis*. *Plant Cell*, 23, 534–549.
- [9] Cabello JV, Arce AL, Chan RL. (2012). The homologous HD-Zip I transcription factors HaHB1 and AtHB13 confer cold tolerance via the induction of pathogenesis-related and glucanase proteins. *Plant J* 69: 141–153.

- [10] Cutler S. R., Rodriguez P. L., Finkelstein RR, Abrams SR. (2010). Abscisic acid: emergence of a core signaling network. *Annual Review of Plant Biology* 61: 651–679.
- [11] Dai, M., Xue, Q., McCray, T., Margavage, K., Chen, F., Lee, J., Nezames, C., Guo, L., Terzaghi, W., Wan, J., Deng, X., Wang, H. (2013). The PP6 phosphatase regulates ABI5 phosphorylation and abscisic acid signaling in *Arabidopsis*. *Plant Cell* 25: 517-34.
- [12] Debeaujon, I., Leon-Kloosterziel, K. M., and Koornneef, M. (2000) Influence of the testa on seed dormancy, germination and longevity in *Arabidopsis thaliana*. *Plant Physiol.* 122: 403-414.
- [13] Dekkers, B. J., Costa, M. C. D., Maia, J., Bentsink, L., Ligterink, W., & Hilhorst, H. W. (2015). Acquisition and loss of desiccation tolerance in seeds: from experimental model to biological relevance. *Planta*, 1-15.
- [14] Finkelstein R. (2013). Abscisic Acid Synthesis and Response. *The Arabidopsis book* 1: 1-36.
- [15] Gonzalez-Guzman, M., Pizzio, G. A., Antoni, R., Vera-Sirera, F., Merilo, E., Bassel, G. W.,... & Rodriguez, P. L. (2012). *Arabidopsis* PYR/PYL/RCAR receptors play a major role in quantitative regulation of stomatal aperture and transcriptional response to abscisic acid. *The Plant Cell Online*, 24 (6), 2483-2496.
- [16] Park, S. Y., Fung, P., Nishimura, N., Jensen, D. R., Fujii, H., Zhao, Y., & Cutler, S. R. (2009). Abscisic acid inhibits type 2C protein phosphatases via the PYR/PYL family of START proteins. *Science*, 324 (5930), 1068-1071.
- [17] Hoekstra, F. A., Golovina, E. A., & Buitink, J. (2001). Mechanisms of plant desiccation tolerance. *Trends in plant science*, 6 (9), 431-438. <http://www.arabidopsis.org>
- [18] Huijser, C., Kortstee, A., PEGo, J., Weisbeek, P., Wisman, E., & Smeeckens, S. (2000). The *Arabidopsis* SUCROSE UNCOUPLED - 6 gene is identical to ABCISIC ACID INSENSITIVE - 4: involvement of abscisic acid in sugar responses. *The Plant Journal*, 23 (5), 577-585.
- [19] Iuchi S., Kobayashi M., Yamaguchi-Shinozaki K. & Shinozaki K. (2000). A stress-inducible gene for 9-cis-epoxycarotenoid dioxygenase involved in abscisic acid biosynthesis under water stress in drought-tolerant cowpea. *Plant Physiology* 123, 553–562.
- [20] Komatsu, K., Suzuki, N., Kuwamura, M., Nishikawa, Y., Nakatani, M., Ohtawa, H., & Sakata, Y. (2013). Group APP2Cs evolved in land plants as key regulators of intrinsic desiccation tolerance. *Nature communications*, 4.
- [21] Krasensky, J., & Jonak, C. (2012). Drought, salt, and temperature stress-induced metabolic rearrangements and regulatory networks. *Journal of Experimental Botany*, 63 (4), 1593-1608.
- [22] Krysan, P. J., Young, J. C., & Sussman, M. R. (1999). T-DNA as an insertional mutagen in *Arabidopsis*. *The Plant Cell Online*, 11 (12), 2283-2290.
- [23] Lee, K. H., Piao, H. L., Kim, H. Y., Choi, S. M., Jiang, F., Hartung, W., & Hwang, I. (2006). Activation of glucosidase via stress-induced polymerization rapidly increases active pools of abscisic acid. *Cell*, 126 (6), 1109-1120.
- [24] Lim, E.-K., Doucet, C. J., Hou, B., Jackson, R. G., Abrams, S. R., Bowles, D. J. (2005). Resolution of (+)-abscisic acid using an *Arabidopsis* glycosyltransferase. *Tetrahedron: Asymmetry* 16: 143-147.
- [25] Lopez-Molina, L., Chua, N.-H. (2000). A null mutation in a bZIP factor confers ABA-insensitivity in *Arabidopsis thaliana*. *Plant Cell Physiol.* 41: 541-547.
- [26] Lopez-Molina, L., Mongrand, S., Chua, N.-H. (2001). A post-germination developmental arrest checkpoint is mediated by abscisic acid and requires the ABI5 transcription factor in *Arabidopsis*. *Proc. Natl. Acad. Sci. USA* 98: 4782-4787. Ma Y, Szostkiewicz I, Korte A, Moes D, Yang Y, Christmann A, Grill E. 2009. Regulators of PP2C phosphatase activity function as abscisic acid sensors. *Science* 324: 1064–1068.
- [27] Maia, J., Dekkers, B. J., Dolle, M. J., Ligterink, W., & Hilhorst, H. W. (2014). Abscisic acid (ABA) sensitivity regulates desiccation tolerance in germinated *Arabidopsis* seeds. *New Phytologist*, 203 (1), 81-93.
- [28] Maia J, Dekkers BJW, Provart NJ, Ligterink W, Hilhorst HWM. (2011). The re-establishment of desiccation tolerance in germinated *Arabidopsis thaliana* seeds and its associated transcriptome. *PLoS ONE* 6: e29123.
- [29] Milla, M. A. R., Townsend, J., Chang, F., & Cushman, J. C. (2006). The *Arabidopsis* AtDi19 gene family encodes a novel type of Cys2/His2 zinc-finger protein implicated in ABA-independent dehydration, high-salinity stress and light signaling pathways. *Plant molecular biology*, 61 (1-2), 13-30.
- [30] Miyakawa T, Fujita Y, Yamaguchi-Shinozaki K, Tanokura M. (2013). Structure and function of abscisic acid receptors. *Trends in Plant Science* 18: 259–266.
- [31] Munnik, T. and Vermeer, J. E. (2010). Osmotic stress-induced phosphoinositide and inositol phosphate signaling in plants. *Plant Cell Environ.* 33, 655–669.
- [32] Okamoto, M., Kuwahara, A., Seo, M., Kushiro, T., Asami, T., Hirai, N., Kamiya, Y., Koshihara, T., Nambara, E. (2006). CYP707A1 and CYP707A2, which encode abscisic acid 8'-hydroxylases, are indispensable for proper control of seed dormancy and germination in *Arabidopsis*. *PlantPhysiol.* 141: 97-107.
- [33] Ribone PA, Capella M, Chan RL. (2015) Functional characterization of the homeodomain leucine zipper I transcription factor AtHB13 reveals a crucial role in *Arabidopsis* development. *J Exp Bot* 66: 5929–5943.
- [34] Santiago, J., Dupeux, F., Round, A., Antoni, R., Park, S. Y., Jamin, M.,... & Márquez, J. A. (2009). The abscisic acid receptor PYR1 in complex with abscisic acid. *Nature*, 462 (7273), 665-668.
- [35] Schwartz, S. H., & Zeevaart, J. A. (2010). Abscisic acid biosynthesis and metabolism. *Plant Hormones*, 137-155.
- [36] Schwartz, S. H., Qin, X., & Zeevaart, J. A. (2003). Elucidation of the indirect pathway of abscisic acid biosynthesis by mutants, genes, and enzymes. *Plant Physiology*, 131 (4), 1591-1601.
- [37] Shinozaki, K., & Yamaguchi-Shinozaki, K. (2007). Gene networks involved in drought stress response and tolerance. *Journal of experimental botany*, 58 (2), 221-227.
- [38] Shinozaki K., Yamaguchi-Shinozaki K. & Seki M. (2003). Regulator network of gene expression in the drought and cold stress responses. *Current Opinion in Plant Biology* 6, 410–417.

- [39] Sun, Q., Csorba, T., Skourti-Stathaki, K., Proudfoot, N. J., & Dean, C. (2013). R-loop stabilization represses antisense transcription at the Arabidopsis FLC locus. *Science*, 340 (6132), 619-621.
- [40] Taiz, L., & Zeiger, E. (2006). *Stress physiology*. Plant physiology, 4.
- [41] Tan B. C., Joseph L. M., Deng W. T., Liu L., Li Q. B., Cline K. & McCarty D. R. (2003). Molecular characterization of the Arabidopsis 9-cis epoxycarotenoid dioxygenase gene family. *Plant Journal* 35, 44–56.
- [42] Tran, L.-S. P., Nakashima, K., Sakuma, Y., Simpson, S. D., Fujita, Y., Maruyama, K., Fujita, M., Seki, M., Shinozaki, K., Yamaguchi-Shinozaki, K. (2004). Isolation and functional analysis of Arabidopsis stress inducible NAC transcription factors that bind to a drought-responsive cis-element in the early responsive to dehydration stress 1 promoter. *Plant Cell* 16: 2481-2498.
- [43] Tripathi, P., Rabara, R. C., & Rushton, P. J. (2014). A systems biology perspective on the role of WRKY transcription factors in drought responses in plants. *Planta*, 239 (2), 255-266.
- [44] Umezawa, T., Okamoto, M., Kushiro, T., Nambara, E., Oono, Y., Seki, M.,... & Shinozaki, K. (2006). CYP707A3, a major ABA 8' - hydroxylase involved in dehydration and rehydration response in Arabidopsis thaliana. *The Plant Journal*, 46 (2), 171-182.
- [45] Umezawa, T., Nakashima, K., Miyakawa, T., Kuromori, T., Tanokura, M., Shinozaki, K., & Yamaguchi-Shinozaki, K. (2010). Molecular basis of the core regulatory network in ABA responses: sensing, signaling and transport. *Plant and cell physiology*, 51 (11), 1821-1839.
- [46] Verdier, J., Lalanne, D., Pelletier, S., Torres-Jerez, I., Righetti, K., Bandyopadhyay, K., & Buitink, J. (2013). A regulatory network-based approach dissects late maturation processes related to the acquisition of desiccation tolerance and longevity of Medicago truncatula seeds. *Plant physiology*, 163 (2), 757-774.
- [47] Westwood, J. H., Mccann, L., Naish, M., Dixon, H., Murphy, A. M., Stancombe, M. A., & Carr, J. P. (2013). A viral RNA silencing suppressor interferes with abscisic acid - mediated signaling and induces drought tolerance in Arabidopsis thaliana. *Molecular plant pathology*, 14 (2), 158-170.
- [48] Wind, J. J., Peviani, A., Snel, B., Hanson, J., & Smeekens, S. C. (2013). ABI4: versatile activator and repressor. *Trends in plant science*, 18 (3), 125-132.
- [49] Xiong, L., & Zhu, J. K. (2003). Regulation of abscisic acid biosynthesis. *Plant physiology*, 133 (1), 29-36.
- [50] Xiong, L., Schumaker, K. S., & Zhu, J. K. (2002). Cell signaling during cold, drought, and salt stress. *The Plant Cell Online*, 14 (suppl 1), S165-S183.
- [51] Yamaguchi-Shinozaki, K., & Shinozaki, K. (2005). Organization of cis-acting regulatory elements in osmotic- and cold-stress-responsive promoters. *Trends in plant science*, 10 (2), 88-94.
- [52] Yang, C., Li, D., Mao, D., Liu, X., Ji, C., Li, X.,... & Zhu, L. (2013). Overexpression of microRNA319 impacts leaf morphogenesis and leads to enhanced cold tolerance in rice (*Oryza sativa* L.). *Plant, cell & environment*, 36 (12), 2207-2218.
- [53] Ye, N., Zhu, G., Liu, Y., Li, Y., & Zhang, J. (2011). ABA controls H₂O₂ accumulation through the induction of OsCATB in rice leaves under water stress. *Plant and cell physiology*, 52 (4), 689-698.