

# Kinetics and Mechanism of Electron-Transfer Reactions: Oxidation of Nalidixic Acid by Diperiodatocuprate (III) in Alkaline Medium

Gajala Tazwar, Mahima Sharma, Vijay Devra<sup>\*</sup>

Department of Chemistry, Janki Devi Bajaj Government Girls College, Kota, India

## **Email address**

v\_devra1@rediffmail.com (V. Devra) \*Corresponding author

# Citation

Gajala Tazwar, Mahima Sharma, Vijay Devra. Kinetics and Mechanism of Electron-Transfer Reactions: Oxidation of Nalidixic Acid by Diperiodatocuprate (III) in Alkaline Medium. *World Journal of Biochemistry and Molecular Biology*. Vol. 3, No. 2, 2018, pp. 37-45.

Received: February 27, 2018; Accepted: March 21, 2018; Published: May 16, 2018

**Abstract:** The kinetics and mechanism of oxidation of nalidixic acid by diperiodatocuprate (III) in aqueous alkaline medium has been studied spectrophotometrically at 303 K. The reaction exhibits first order with respect to oxidant but substrate dependence is complex. The stoichiometry of the reaction has been observed to two moles of the oxidant for a mole of the substrate. The oxidation product of the substrate is 1-ethyl-2-hydroxy-1, 4-dihydro-7-methyl-4-oxo-1, 8-naphthyridine-3-carboxylic acid as established spectrally. The active species of diperiodatocuprate (III) is understood to be as monoperiodatocuprate (III). The activation parameters were also determined and discussed. The activation parameters and thermodynamics quantities were also determined and discussed. A plausible reaction mechanism has been suggested to account for experimental observations.

Keywords: Diperiodatocuprate (III), Nalidixic Acid, Oxidation, Kinetics, Mechanism

# 1. Introduction

In recent years, the study of highest oxidation state of transition metals has frame-upped by many researchers. Transition metals in a higher oxidation state can be stabilized by chelation with suitable polydentate ligands. Metal chelate with such as diperiodatocuprate (III), diperiodatoargenate (III) and diperiodatonickelate (IV) are good oxidants [1-3]. Diperiodatocuprate (III) (DPC) is a versatile one-electron oxidant for various organic compounds in alkaline medium and its use as an analytical reagent is now well recognized [4]. Copper complexes have engaged a major place in oxidation chemistry due to their prosperity and applicability in biological chemistry [5]. The study using DPC is novel and confined to few cases owing to its limited solubility and stability [6-13].

Fluoroquinolones are a family of synthetic antibacterial agents with a rising popularity. These antibiotics display a broad spectrum of antibacterial activity including strong effects on gram-negative aerobic and anaerobic organisms as well as on gram-positive and a typical pathogens [14, 15]. But these are not fully metabolized in the body and are partially excreted in its pharmaceutically active form [16, 17].

Due to the limited biodegradability and widespread use of these antibiotics, an incomplete removal is obtained in typical waste water treatment plants and analogous huge quantities are discharged into the environment. As a result, numerous antibiotics can be found in surface waters producing injurious effects on aquatic organisms [18, 19]. For the removal of these fluoroquinolone many studies are used, in which oxidation process is mostly used and degrade them from the environment [20, 21]. Wang et al have select ClO<sub>2</sub> as an oxidant to understand degradation of drugs in possible products [22]. Zhang et al used MnO<sub>2</sub> as oxidant in drug oxidation for the elucidation of their oxidant product [23]. However Nalidixic acid (NA) was not used in their studies, neither in quantification nor in the kinetic studies of oxidative degradation with DPC. Nalidixic acid (NA) with molecular formula C12H12N2O3 (1-ethydm-3, 4-dihydro-7methyl-4-oxo-1, 8-naphthyridine-3-carboxylic acid) is the first synthesized antimicrobial quinolone. NA is an ionizable, non-biodegradable photosensitive molecule [24, 25] with a carboxylic acid function having a pKa of 5.95 [26]. NA is an antibacterial drug still widely used for urinary tract infections [27]. However the kinetics of oxidation of medicinally

important compound such as NA with the following view point: first, identify the active species of DPC in aqueous alkaline medium, secondly, whether or not the substrate forms any intermediate complex before undergoing to final reaction products. Hence, the present investigation is aimed to elucidate the reactivity of NA towards DPC, to arrive at a plausible mechanism and to understand the reactive species.

# **2.** Materials and Methods

## 2.1. Chemicals

All chemicals used were of analytical grade and standard solution of NA (KORES India Limited) was prepared by dissolving calculating quantity of pure drug in double distilled water. A stock solution of periodate was prepared by dissolving known amount of  $KIO_4$  (MERCK) in hot water and used after 24 h and concentration was determined iodometrically [28]. The complex of copper (III) periodate was prepared [29] and standardized by a known method [30]. KOH and KNO<sub>3</sub>(BDH) were employed to maintain the required alkalinity and ionic strength respectively in reaction solutions. Doubly distilled water was employed throughout the study.

#### **2.2. Instrumentation**

For kinetic study, a Peltier accessory (temperature-Controlled) attached to a double beam U.V. 3000<sup>+</sup>, UV-Visible spectrophotometer (LABINDIA) with U.V. path length 1.0 cm in the spectral range 200-800 nm, was used. Liquid Chromatography-Mass Spectroscopy (LC-MS), (Q-TOF Micromass, WATERS Company, UK) and Fourier Transform Infrared (FT-IR) Spectrophotometer (ALPHA-T, Bruker, Germany) were used for product analysis.

#### **2.3. Kinetic Procedure**

All reaction ingredients except DPC were taken in flask and the reaction was initiated by adding requisite volume of temperature pre-equilibrated solution of DPC. The progress of the reaction was followed by measuring the absorbance of unreacted DPC in the reaction mixture by UV-Visible spectrophotometer at 415 nm. The application of Beer's law of DPC at 415 nm has been verified. The molar absorptivity index of DPC was found to be  $\varepsilon = 6230 \text{ dm}^3 \text{mol}^{-1} \text{cm}^{-1}$  [31]. Since kinetics was studied under pseudo first order conditions ([NA])>>10[DPC]). Pseudo first order plots were obtained and pseudo-first-order rate constants ( $k_{obs}$ ) were calculated from these plots. The value of  $k_{obs}$  was reproducible within ±5%.

# **3. Results and Discussion**

#### **3.1. Stoichiometry and Product Analysis**

The stoichiometry of the reaction was calculated by taken excess of DPC over NA in a flask thermostated at  $30\pm0.1^{\circ}$ C. The estimation of excess DPC corresponds to the stoichiometry of the reaction as represented by the equation (1).



Figure 1. LC-MS spectra of oxidation product of NA.



Figure 2. FT-IR spectra of the product of oxidation of NA by DPC.

The main oxidation product of NA 1-ethyl-2-hydroxy-1, 4dihydro-7-methyl-4-oxo-1, 8-naphthyridine-3-carboxylic acid was isolated with the help of TLC and characterized by LC-MS, FT-IR spectral analysis. LC/MS analysis of the reaction indicated the presence of a product with molecular ion of m/z 248 corresponds to 1-ethyl-2-hydroxy-1, 4dihydro-7-methyl-4-oxo-1, 8-naphthyridine-3-carboxylic acid (Figure 1). The molecular ion of nalidixic acid is m/z 232.2. The IR spectroscopy shows a broad peak at 3382.39 cm<sup>-1</sup> which is due to -OH stretching (Figure 2) and the remaining peaks are of the parent compound.

#### 3.2. Diperiodatocuparate (III) Dependence

The concentration of DPC was varied from  $1.0 \times 10^{-5}$  to  $10.0 \times 10^{-5}$  mol dm<sup>-3</sup> at [NA] =  $5.0 \times 10^{-4}$  mol dm<sup>-3</sup>, alkali = 0.05 mol dm<sup>-3</sup> and I = 0.10 mol dm<sup>-3</sup> at 30°C temperature. The plot of log absorbance versus time was linear (Figure 3) indicating the reaction is first order with respect to [DPC]. The observed pseudo first order rate constant ( $k_{obs}$ ) were independent of the initial concentration of DPC (Table 1).



*Figure 3.* First order plots of the variation of DPC concentration at 30°C.

Conditions:  $[NA] = 5.0 \times 10^{-4}$ ,  $[OH^{-}] = 5.0 \times 10^{-2}$ ,  $[IO_{4}^{-}] = 5.0 \times 10^{-5}$  and I = 0.1 /mol dm<sup>-3</sup>.  $[DPC] \times 10^{-5}$  mol dm<sup>-3</sup> = (A) 1.0, (B) 2.0, (C) 3.0, (D) 4.0, (E) 5.0, (F) 6.0, (G) 7.5, (H) 10.0.

#### 3.3. Nalidixic acid Dependence

The concentration of NA was varied from  $1.0 \times 10^{-4}$  to  $10.0 \times 10^{-4}$  mol dm<sup>-3</sup> at constant concentration of DPC, alkali, IO<sub>4</sub><sup>-</sup> and ionic strength at 25°C, 30°C and 35°C respectively. The rate of reaction increases with increasing concentration of NA initially but levels off at higher concentration indicating complex order with respect to NA (Table 1). This was confirmed by the plot of 1/k<sub>obs</sub> versus 1/[NA] which was linear with a positive intercept.

<b>Table 1.</b> First order rate constants $(k_{obs})$ for the reaction of NA with DPC	in
aqueous alkaline medium at $I = 0.1$ mol dm <sup>-3</sup> and Temperature = 303 K.	

10 <sup>5</sup> [DPC]	10 <sup>4</sup> [NA]	10 <sup>2</sup> [OH <sup>-</sup> ]	10 <sup>4</sup> [KIO <sub>4</sub> ]	10 <sup>4</sup> k <sub>obs</sub>
(mol dm <sup>-3</sup> )	(mol dm <sup>-3</sup> )	(mol dm <sup>-3</sup> )	(mol dm <sup>-3</sup> )	(s <sup>-1</sup> )
1.0	5.0	5.0	5.0	3.16
2.0	5.0	5.0	5.0	3.12
3.0	5.0	5.0	5.0	3.08
4.0	5.0	5.0	5.0	3.18
5.0	5.0	5.0	5.0	3.15
6.0	5.0	5.0	5.0	3.12
7.5	5.0	5.0	5.0	3.05
10.0	5.0	5.0	5.0	3.15
5.0	1.0	5.0	5.0	0.99
5.0	2.0	5.0	5.0	1.71
5.0	3.0	5.0	5.0	2.32
5.0	4.0	5.0	5.0	2.81
5.0	5.0	5.0	5.0	3.15
5.0	6.0	5.0	5.0	3.43
5.0	7.5	5.0	5.0	3.72
5.0	10.0	5.0	5.0	3.91
5.0	5.0	2.0	5.0	1.42
5.0	5.0	3.0	5.0	2.01
5.0	5.0	4.0	5.0	2.65
5.0	5.0	5.0	5.0	3.15
5.0	5.0	6.0	5.0	3.45
5.0	5.0	7.5	5.0	3.92
5.0	5.0	10.0	5.0	4.50
5.0	5.0	5.0	1.0	6.85
5.0	5.0	5.0	2.5	5.05
5.0	5.0	5.0	5.0	3.15
5.0	5.0	5.0	7.5	2.35
5.0	5.0	5.0	10.0	1.97

## **3.4. Hydroxyl ion Dependence**

Hydroxyl ion concentration was varied from  $2.0 \times 10^{-2}$  to  $10.0 \times 10^{-2}$  mol dm<sup>-3</sup> at fixed concentration of other reaction ingredients [DPC] =  $5.0 \times 10^{-5}$ , [NA] =  $5.0 \times 10^{-4}$ , [IO<sub>4</sub><sup>-</sup>] =  $2.0 \times 10^{-5}$ , I = 0.1 / mol dm<sup>-3</sup> at three temperature 25°C,  $30^{\circ}$ C,  $35^{\circ}$ C respectively. Pseudo first order rate constant ( $k_{obs}$ ) was found to be increase with increase in [OH<sup>-</sup>] (Table 1). A plot of log  $k_{obs}$  versus log [OH<sup>-</sup>] was linear with a fractional slope of 0.56. This was confirmed by the plot of  $1/k_{obs}$  versus 1/[OH<sup>-</sup>].

#### 3.5. Periodate Ion Dependence

The effect of concentration of  $\text{KIO}_4$  was studied by different periodate concentration from  $1.0 \times 10^{-5}$  to  $1.0 \times 10^{-4}$  mol dm<sup>-3</sup>, keeping other reactant concentrations constant. It was found added periodate had a retarding effect on the rate of reaction (Table 1). The order with respect to periodate concentration was found to be less than unity with negative slope.

#### 3.6. Effect of Ionic Strength and Dielectric Constant

At constant concentration of reactants and other conditions, the ionic strength was varied by different concentration of sodium nitrate from 0.20 to 0.50 mol dm<sup>-3</sup>. The rate of reaction is unaffected by variations of ionic strength. The effect of the dielectric constant (D) was studied by different t-butanol-water content (v/v) in the reaction mixture with all other conditions maintained constant. The dielectric constant of the reaction medium has no effect on the rate of reaction.

### **3.7. Test for Free Radical**

Acrylonitrile was added in the reaction mixture and kept in an inert atmosphere ten diluted with methanol. White precipitate comes, indicating the formation of free radicals during reaction.

#### 3.8. Mechanism

The present investigation carried out in aqueous alkaline medium so involvement of periodate in multiple equilibria (2-4) depending upon the pH of solution [32].

$$H_5 IO_6 \rightleftharpoons H_4 IO_6^- + H^+ \tag{2}$$

$$H_4 IO_6^- \rightleftharpoons H_3 IO_6^{2-} + H^+ \tag{3}$$

$$H_3 IO_6^{2-} \rightleftharpoons H_2 IO_6^{3-} + H^+ \tag{4}$$

In acid medium periodate acid exist as  $H_5IO_6^-$  and at pH = 7 as  $H_4IO_6^-$ . In this study employed alkaline medium, the main species are expected to be  $H_3IO_6^{2-}$  and  $H_2IO_6^{3-}$ . Thus soluble copper (III) periodate complex exist as  $[Cu(OH)_2(H_3IO_6)_2]^{3-}$  [33].

The stiochiometry of the reaction between diperiodatocuprate (III) complex and NA is an alkaline medium has 1:2. The rate of reaction increases with alkali concentration can be explained by formation of  $[Cu(OH)_2(H_3IO_6)]^{4-}$  from  $[Cu(OH)_2(H_3IO_6)_2]^{3-}$ .

$$[Cu(OH)_2(H_3IO_6)_2]^{3-} + OH^- \underbrace{\overset{K_1}{\longleftarrow}} [Cu(OH)_2(H_3IO_6)(H_2IO_6)]^{4-} + H_2O$$
(5)

The rate of reaction decrease with increase in periodate concentration suggests that equilibrium of Cu (II) periodate complex

to form monoperiodato cuprate (III) (MPC) [31].

$$[Cu(OH)_2(H_3IO_6)(H_2IO_6)]^{4-} \xrightarrow{K_2} [Cu(OH)_2(H_3IO_6)]^{-} + (H_2IO_6)^{3-}$$
(6)

It can also possible that [MPC] is more important than [DPC] in the reaction.

The experimental results shows fractional order with respect to NA concentration indicates complex formation with NA and DPC species [34] before the rate determining step. The braking of this complex (C) is assigned as slowest step, leading to formation of NA radical intermediate. The radical intermediate reacts with other molecules of MPC species, to give final products (Scheme 1). The effect of ionic strength on the rate explains qualitatively the involvement of neutral molecule in the reacts on the basis of results, mechanism are as follows.

$$Cu(HL)_{2} + [OH^{-}] \rightleftharpoons CuL(HL) + H_{2}O$$

$$CuL(HL) \rightleftharpoons Cu(HL) + L$$

$$Cu(HL) + S \rightleftharpoons Complex(C)$$

$$Complex(C) \rightarrow S^{*} + Cu(II)$$

$$S^{*} + Cu(HL) \rightarrow \operatorname{Pr} oducts + Cu(II)$$

So, the detailed mechanistic scheme for the oxidation of nalidixic acid by DPC is as follows:



Free radical + 
$$[Cu(OH)_2(H_3IO_6)]^-$$
 +  $OH^-$  Fast  
H<sub>3</sub>C N N OH +  $Cu(OH)_2$  +  $H_3IO_6^{2-}$  (9)

Scheme 1. Proposed mechanism for the oxidation of NA by DPC in aqueous alkaline medium. From the above mechanism the following rate law (10) - (20) can be derived.

$$Rate = -\frac{d[DPC]}{dt} = k[C] = \frac{kK_1K_2K_3[Cu(OH)_2(H_3IO_6)]^{-}[NA][OH^{-}]}{[H_3IO_6^{3^{-}}]}$$
(10)

The total concentration of DPC is given by equation (11)

$$[DPC]_{t} = [DPC]_{f} + [Cu(OH)_{2}(H_{3}IO_{6})(H_{2}IO_{6})]^{4-} + [Cu(OH)_{2}(H_{3}IO_{6})]^{-} + [Complex(C)]$$
(11)

$$[DPC]_{t} = [DPC]_{f} + K_{1}[DPC][OH^{-}] + \frac{K_{1}K_{2}[DPC][OH^{-}]}{[H_{2}IO_{6}^{3-}]} + \frac{K_{1}K_{2}[DPC][NA][OH^{-}]}{[H_{2}IO_{6}^{3-}]}$$

$$[DPC]_{t} = [DPC]_{f} \left\{ \frac{[H_{2}IO_{6}^{3-}] + K_{1}[H_{2}IO_{6}^{3-}][OH^{-}] + K_{1}K_{2}[OH^{-}] + K_{1}K_{2}[DPC][NA][OH^{-}]}{[H_{2}IO_{6}^{3-}]} \right\}$$
(12)

$$[DPC]_{f} = \frac{[1 - V_{1}f_{1} - V_{2} - V_{6}]}{[H_{2}IO_{6}^{3-}] + K_{1}[H_{2}IO_{6}^{3-}][OH^{-}] + K_{1}K_{2}[OH^{-}] + K_{1}K_{2}[DPC][NA][OH^{-}]}$$
(13)

Where [DPC]t and [DPC]f presents total and free concentrations of DPC respectively. Similarly total concentration of Nalidixic acid is given by

$$[NA]_{t} = [NA]_{f} + [C]$$

$$[NA]_{t} = [NA]_{f} + \frac{K_{1}K_{2}K_{3}[DPC]^{-}[NA][OH^{-}]}{[H_{3}IO_{6}^{3-}]}$$

$$[NA]_{t} = \frac{[NA]_{f}[H_{3}IO_{6}^{3-}] + K_{1}K_{2}K_{3}[DPC]^{-}[OH^{-}]}{[H_{3}IO_{6}^{3-}]}$$

$$[NA]_{f} = \frac{[NA]_{t}[H_{3}IO_{6}^{3-}]}{[H_{3}IO_{6}^{3-}] + K_{1}K_{2}K_{3}[DPC]^{-}[OH^{-}]}$$
(14)

At low concentration of DPC, OH<sup>-</sup> and H<sub>2</sub>IO<sub>6</sub><sup>-</sup> used.

$$[NA]_f = [NA]_t \tag{15}$$

Similarly,

$$[OH^{-}]_{t} = [OH]_{f} + [Cu(OH)_{2}(H_{3}IO_{6})(H_{2}IO_{6})]^{4-} + [Cu(OH)_{2}(H_{3}IO_{6})]^{-} + [Complex(C)]$$

 $[OH^{-}]_{t} = [OH]_{f} + K_{1}[OH^{-}][Cu(OH)_{2}(H_{3}IO_{6})_{2}]^{-} + \frac{K_{1}K_{2}[Cu(OH)_{2}(H_{3}IO_{6})_{2}]^{3^{-}}[OH^{-}]}{[H_{2}IO_{6}^{3^{-}}]} + \frac{K_{1}K_{2}[Cu(OH)_{2}(H_{3}IO_{6})_{2}]^{3^{-}}[NA]}{[H_{2}IO_{6}^{3^{-}}]}$ (16)

At low concentration of DPC, [NA] and H<sub>2</sub>IO<sub>6</sub><sup>3-</sup> used,

$$[OH^-]_f = [OH^-]_t \tag{17}$$

Substituting equations (13),(15),(17) in equation (10) and omitting 't' and 'f' subscripts.

$$Rate = -\frac{d[DPC]}{dt} = \frac{kK_1K_2K_3[DPC][NA][OH^-]}{[H_2IO_6^{3-}] + K_1[H_2IO_6^{3-}][OH^-] + K_1K_2[OH^-] + kK_1K_2[NA][OH^-]}$$
(18)

or

$$k_{obs} = -\frac{Rate}{[DPC]} = \frac{kK_1K_2K_3[NA][OH^-]}{[H_2IO_6^{3-}] + K_1[H_2IO_6^{3-}][OH^-] + K_1K_2[OH^-] + kK_1K_2[NA][OH^-]}$$
(19)

The equation (19) can be rearranged as

$$\frac{1}{k_{obs}} = \frac{[H_2 I O_6^{3-}]}{k K_1 K_2 K_3 [NA] [OH^-]} + \frac{[H_2 I O_6^{3-}]}{k K_2 K_3 [NA]} + \frac{1}{k K_3 [NA]} + \frac{1}{k}$$
(20)

According to equation (20) the plot of  $1/k_{obs}$  versus  $1/[OH^-]$ ,  $1/k_{obs}$  versus 1/[NA] and  $1/k_{obs}$  versus  $[H_2IO_6^{3-}]$  are linear

The thermodynamics quantities of the first, second and third equation steps of scheme 1 were calculated by variation of  $[H_2IO_6^{3-}]$ , [NA],  $[OH^-]$  at three temperatures (Table 1). The plots of  $1/k_{obs}$  versus  $1/[OH^-]$ ,  $1/k_{obs}$  versus 1/[NA] and  $1/k_{obs}$  versus  $[H_2IO_6^{3-}]$  are obtained linear (Figure 4, 5, 6)

respectively). From the slopes and intercept, the values of  $K_1$  were calculated at three temperatures (Table 2). A Von't Haff plot was made by log  $K_1$  versus 1/T and values of enthalpy of reaction ( $\Delta H$ ), entropy of reaction ( $\Delta S$ ) and free energy of reaction ( $\Delta G$ ) were calculated. A comparison of latter values with those obtained for the slow step of reaction, clearly indicates the reaction before rate determining step is fast with law activation energy [35]. In the similar manner the  $K_2$  and  $K_3$  values were calculated and thermodynamics quantities are given in table 2.



 $[DPC] = 5.0 \times 10^{-5}$ ,  $[NA] = 5.0 \times 10^{-4}$ ,  $[IO_4^-] = 5.0 \times 10^{-4}$  and I = 0.1 /mol dm<sup>-3</sup>. Figure 4. Plot of  $1/k_{obs}$  versus  $1/[OH^-]$  at 30 °C.







 $[DPC] = 5.0 \times 10^{-5}$ ,  $[NA] = 5.0 \times 10^{-4}$ ,  $[OH^{-1}] = 5.0 \times 10^{-2}$  and I = 0.1 /mol dm<sup>-3</sup>.

Figure 6. Plot of  $1/k_{obs}$  versus  $[H_2IO_6^{3-}]$  at 30 °C.

The negative value of  $\Delta S^{\#}$  indicates complex is more ordered than reactants [35]. The value of  $\Delta H^{\#}$  and  $\Delta S^{\#}$  were favourable for electron transfer reactions. The value of

entropy of activation and law value of activation and higher value of rate constant of slow step indicates the oxidation reaction occurs via inner-sphere mechanism [36].

Table 2. Activation parameters and thermodynamics quantities for the oxidation of nalidixic acid by DPC in aqueous alkaline medium.

Temperature (K)	$10^3 \mathrm{k}(\mathrm{s}^{-1})$	Activation parameters	
298	5.88	Ea (kJ mol <sup>-1</sup> )	19.47
303	6.58	$\Delta H^{\#}$ (kJ mol <sup>-1</sup> )	16.95
308	7.58	$\Delta S^{\#} (J K^{-1} mol^{-1})$ $\Delta G^{\#} (kJmol^{-1})$	-183.43 86.91
Temperature (K)	$K_1 (mol^{-1} dm^3)$	$10^2 \text{ K}_2 \text{ (mol}^{-1} \text{ dm}^3)$	$K_3 10^5 \text{ (mol dm}^{-3}\text{)}$
298	8.17	15.53	19.11
303	7.04	17.78	25.10
308	5.78	20.72	29.70
Thermodynamic quantities	Values from K <sub>1</sub>	Values from K <sub>2</sub>	Values from K <sub>3</sub>
$\Delta H (kJ mol^{-1})$	-26.29	21.92	22.52
$\Delta S (J K^{-1} mol^{-1})$	-315.54	-187.07	-48.25
$\Delta G (kJmol^{-1})$	69.16	78.60	37.06

# 4. Conclusion

Among the various species of copper (III) in alkaline medium, monoperiodatocuprate (III),  $[Cu(H_2IO_6)(H_2O)_2]$  is consider to be active species for the title reaction. The results indicate that, the rate of pH in the reaction medium is important. Rate constant of the slow step and other equation constants involved in the mechanism evaluated and activation parameters with respect to slow step indicates the oxidation reaction occur via inner-sphere mechanism.

## References

 H. Yao, M. Zhang, W. Zeng, X. Zeng, Z. Zhang, A novel chemiluminescence assay of mitoxantrone based on diperiodatocuprate (III) oxidation. Spectrochim. Acta. Part A. 117 (2014) 645–650.

- [2] A. Kumar, P. Kumar, P. Ramamurthy, Kinetics of oxidation of glycine and related substrates by diperiodatoargentate (III). Polyhedron. 18 (1999) 773–780.
- [3] R. S. Shettar, S. T. Nandibewoor, Kinetic, mechanistic and spectral investigations of ruthenium (III)- catalysed oxidation of 4-hydroxycoumarin by alkaline diperiodatonickelate (IV) (stopped flow technique). J. Mol. Catal. A. 234 (2005) 137–143.
- [4] Y. Hu, G. Li, Z. Zhang, A flow injection chemiluminescence method for the determination of lincomycin in serum using a diperiodato-cuprate (III)–luminol system. Luminescence. 26 (2011) 313–318.
- [5] J. E. Weder, C. T. Dillon, T. W. Hambley, B. J. Kennedy, P. A. Lay, J. R. Biffin, H. L. Regtop, N. M. Davies, Copper complexes of non-steroidal anti-inflammatory drugs: an opportunity yet to be realized. Coord. Chem. Rev. 232 (2002) 95–126.
- [6] A. M. Bagoji, P. A. Magdum, S. T. Nandibewoor, Oxidation of Acebutolol by Copper (III) Periodate Complex in Aqueous Alkaline Medium: A Kinetic and Mechanistic Approach. J. Solution Chem. (2016). doi: 10.1007/s10953-016-0539-x

- [7] R. N. Hegde, N. P. Shetti, S. T. Nandibewoor, Oxidative degradation and deamination of atenolol by diperiodatocuprate (III) in aqueous alkaline medium: A mechanistic study, Polyhedron. 28 (2009) 3499–3506.
- [8] K. S. Byadagi, R. V. Hosahalli, S. T. Nandibewoor, S. A. Chimatadar, Oxidation of a Anticholinergic Drug Atropine Sulfate Monohydrate by Alkaline Copper (III) Periodate Complex: A Kinetic and Mechanistic Study. Z. Phys. Chem. 226 (2012) 233–249.
- [9] K. Byadagi, M. Meti, S. Nandibewoor, S. Chimatadar, Catalytic Activity of Palladium (II) and Osmium (VIII) on the Oxidation of Chloramphenicol by Copper (III) Periodate Complex in Aqueous Alkaline Medium: A Comparative Kinetic and Mechanistic Approach. Ind. Eng. Chem. Res. (2013). doi: 10.1021/ie400097n
- [10] S. D. Lamani, P. N. Naik, S. T. Nandibewoor, Spectral and Mechanistic Investigation of the Osmium (VIII) Catalyzed Oxidation of Diclofenac Sodium by the Copper (III) Periodate Complex in Aqueous Alkaline Medium J. Solution Chem. 39 (2010) 1291–1310.
- [11] U. R. Bagwan, A. L. Harihar, S. D. Lamani, I. N. Shaikh, A. B. Teradale, Uncatalyzed Oxidation of Anti Tuberculosis drug, Pyrazinamide by Cu (III) Complex in Aqueous Alkaline Media: A Kinetic Approach. Research Journal of Pharmaceutical, Biological and Chemical Sciences. 8 (2017) 1015-1026.
- [12] M. A. Angadi, S. M. Tuwar, Oxidation of Fursemide by Diperiodatocuprate (III) in Aqueous Alkaline Medium—a Kinetic Study J. Solution Chem. 39 (2010) 165–177.
- [13] S. A. Chimatadar, T. Basavaraj, K. A. Thabaj, S. T. Nandibewoor. Ruthenium (III) catalysed oxidation of gabapentin (neurontin) by diperiodatocuprate (III) in aqueous alkaline medium—A kinetic and mechanistic study. J. Mol. Catal. A: Chem. 267 (2007) 65–71.
- [14] C. M. Oliphant, G. M. Green, Quinolones: a comprehensive review. Clin. Pharmacol. 65 (2002) 455-464.
- [15] R. C. Owens, P. G. Ambrose, Antimicrobial safety: focus on fluoroquinolones. Clin. Infect. Dis. 41 (2005) 144–147.
- [16] A. J. Watkinson, E. J. Murby, S. D. Costanzo, Removal of antibiotics in conventional and advanced wastewater treatment: Implications for environmental discharge and wastewater recycling. Water Res. 41 (2007) 4164–4176.
- [17] H. Stass, Metabolism and Excretion of Moxifloxacin. Drugs. 58 (1999) 231–232.
- [18] D. Calamari, E. Zuccato, S. Castiglioni, R. Bagnati, R. Fanelli, Strategic Survey of Therapeutic Drugs in the Rivers Po and Lambro in Northern Italy. Environ. Sci. Technol. 37 (2003) 1241–1248.
- [19] A. A. Robinson, J. B. Belden, M. J. Lydy, Toxicity of fluoroquinolone antibiotics to aquatic organisms. Environ. Toxicol. Chem. 24 (2005) 423–430.
- [20] J. Fick, H. Soderstrom, R. H. Lindberg, C. Phan, M. Tysklind, D. G. J. Larsson, Contamination of surface, ground, and drinking water from pharmaceutical production. Environ. Toxicol. Chem. 28 (2009) 2522–2527.
- [21] O. A. H. Jones, N. Voulvoulis, J. N. Lester, Potential Ecological and Human Health Risks Associated With the Presence of Pharmaceutically Active Compounds in the Aquatic Environment. Crit. Rev. Toxicol. 34 (2004) 335–350.

- [22] P. Wang, H. Yi-Liang, C. H. Ching-Hua, Water Res. 44 (2010) 5989-5998.
- [23] H. C. Zhang, C. H. Haung, Oxidative transformation of Triclosan and Chlorophene by Manganese Oxides. Environ. Sci. Technol. 37 (2003) 2421-2430.
- [24] G. Mascolo, L. Balest, D. Cassano, G. Laera, A. Lopez, A. Pollice, C. Salerno, Biodegradability of pharmaceutical industrial wastewater and formation of recalcitrant organic compounds during aerobic biological treatment. Bioresour Technol. 101 (2010) 2585-2591.
- [25] L. K. Ge, J. W. Chen, X. X. Wei, S. Y. Zhang, X. L. Qiao, X. Y. Cai, Q. Xie, 474 Aquatic Photochemistry of Fluoroquinolone Antibiotics: Kinetics, Pathways, and Multivariate 475 Effects of Main Water Constituents. Environ. Sci. Technol. 44 (2010) 2400-2405.
- [26] D. L. Ross, C. M. Riley, Aqueous solubility's of some variously substituted quinolone antimicrobials. Int. J. Pharm. 63 (1990) 237-250.
- [27] A. M. Barlow. Nalidixic acid in infections of urinary tract. Br. Med. J. 2 (1963) 1308-1310.
- [28] G. P. Panigrahi, P. K. Misro, Kinetics and mechanism of oxidation of aliphatic ketones by sodium metaperiodate: A comparative study of uncatalysed versus osmium tetraoxidecatalysed oxidation. Indian Journal of Chemistry. 16 (1978) 762–766.
- [29] P. K. Murthy, B. Sethuram, T. N. Rao, Kinetics of oxidation of some alcohols by diperiodatocuprate (iii) in alkaline medium. Zeitschrift fur Physikalische Chemie- Leipzig, 262 (1981) 336–340.
- [30] G. H. Jeffery, J. M. Bassett, J. Mendham, R. C. Denny, Vogels' text book of quantitative chemical analysis (5<sup>th</sup> ed., pp. 371–455). Longman: ELBS (1996).
- [31] N. P. Shetti, S. T. Nandibewoor, Kinetic and mechanistic investigations on oxidation of l-tryptophan by diperiodatocuprate (III) in aqueous alkaline medium. Z. Phys. Chem. 223 (2009) 299-317.
- [32] J. C. Bailar, H. J. Emeleus, S. R. Nyholm, A. F. Trotman Dikenson, Comprehensive inorganic chemistry (Vol. 2, p. 1456). Oxford: Pergamon Press (1975).
- [33] K. Bal Reddy, B. Sethuram, T. Navaneeth Rao, Kinetics of oxidative deamination and decarboxylation of some amino acids by diperiodato cuprate (III) in alkaline medium. Indian Journal of Chemistry A. 20 (1981) 395–397.
- [34] D. C. Hiremath, T. S. Kiran, S. T. Nandibewoor, Oxidation of vanillin by diperiodatocuprate (III) in aqueous alkaline medium: A kinetic and mechanistic study by stopped flow technique. Int. J. Chem. Kinet. 39 (2007) 236–244. http://dx.doi.org/10.1002/(ISSN)1097-4601
- [35] A. Weissberger, E. S. Lewis, (Eds.) Investigation of rates and mechanism of reactions in techniques of chemistry, Wiley Interscience, New York, 1974.
- [36] D. C. Hiremath, K. T. Sirsalmath, S. T. Nandibewoor. Osmium (VIII)/ruthenium (III) catalyzed oxidation of L-lysine by diperiodatocuprate (III) in aqueous alkaline medium: A comparative mechanistic approach by stopped flow technique. Catalysis Letters, 122 (2008) 144–154. http://dx.doi.org/10.1007/s10562-007-9361-9